

Phase A review: Stars and Stellar Populations

Andrea Chiavassa



Outline


- Context, the team and the process
- Sciences cases: Cool stars
- Sciences cases: Dynamical and chemical composition of stellar atmospheres
- Sciences cases: Primitive stars
- Required TRLs

Outline

- Context, the team and the process
- Sciences cases: Cool stars
- Sciences cases: Dynamical and chemical composition of stellar atmospheres
- Sciences cases: Primitive stars
- Required TRLs

The team of Phase A

Science Advisory Committee (SAT): 63 experts in High-Res spectrosc.



The team of Phase A

Working group Stars and
Stellar Populations

C. Allende (IAC, **Spain - chair**), A. Korn (Uppsala University, **Sweden - co-chair**)

V. Adibekyan (IACE, **Portugal**), J.-C. Bouret (LAM, **France**), A. Chiavassa (Lagrange, **France**), N. Christlieb (ZAH, **Germany**),

C. Juul Hansen (Niels Bohr Institute, **Danemark**), O. Kochukhov (Uppsala University, **Sweden**), J. R. de Medeiros, A. Mucciarelli (University of Bologna, **Italy**), E. Niemczura (University of Wroclaw, **Poland**) L. Pasquini (**ESO**), G. Pietrzynski (Polish Academy of Sciences, **Poland**), A. Reiners (University of Göttingen, **Germany**), D. Romano (University of Bologna, **Italy**), R. Schiavon (Liverpool John Moores University, **UK**), S. Sousa (IACE, **Portugal**)

The process

Starting point

Reviewing the science cases in the White paper of Maiolino et al. 2013 (arXiv:1310.3163)

Prioritization of Science Cases

Science Cases to TRL and advising for Technical solutions (Technical Team + ESO)

Estimation of costs and Phase A document

The process

Prioritization of Science Cases

Criteria:

- A. **Scientific impact** (e.g., enabling new discoveries, importance for a broad scientific community, interdisciplinarity, Nobel prize potential)
- B. **Feasibility** and time needed
- C. **Competitiveness** with other instruments

The process

The winners are:

A. **Cool stars** (low mass, brown dwarf, red giant stars)

B. **Primitive stars**

C. Dynamical and chemical **composition of stellar atmospheres**

... + 15 additional science cases ranked lower, including stellar-population studies in the Local Group, via individual stars and extragalactic star clusters, pulsating stars, proto-planetary disks and pre-main sequence stars, asteroseismology in the Local Group, Gravity Darkening, Circumstellar disks, metals in white dwarfs

Outline



Context

- Sciences cases: Cool stars
- Sciences cases: Dynamical and chemical composition of stellar atmospheres
- Sciences cases: Primitive stars
- Required TRLs

Science cases: Cool stars

RSG (Chiavassa et al. 2011)
 $M > 8 M_{\odot}$

AGB (Freytag et al. 2017)
 $M = 1-8 M_{\odot}$





$> 500 R_{\odot}$

$> 200 R_{\odot}$


Brown Dwarf (Freytag et al. 2013)
 $M = 13 M_j$

M Dwarf (Wedemeyer et al. 2013)
 $M = 0.2 M_{\odot}$



$0.1 R_{\odot}$



$0.2 R_{\odot}$



Science cases: Cool stars

RSG (Chiavassa et al. 2011)
 $M > 8 M_{\odot}$

AGB (Freytag et al. 2017)
 $M = 1-8 M_{\odot}$





$> 500 R_{\odot}$

$> 200 R_{\odot}$


Brown Dwarf
(Freytag et al. 2013)
 $M = 13 M_{\text{J}}$

M Dwarf
(Wedemayer et al. 2013)
 $M = 0.2 M_{\odot}$



$0.1 R_{\odot}$

$0.2 R_{\odot}$




HIRES wavelength range

Science cases: Cool stars

At the low temperatures found in the atmospheres, the chemistry becomes complex, and a **myriad of atomic and molecular transitions overlap** in any given part of the spectrum.




This spectral *crowding* causes serious **difficulties to identify and interpret the observations**.

HIRES will observe at or beyond the limit imposed by the thermal and turbulent width of the spectral lines, and in the infrared, where there is a reduced density of transitions.





Science cases: Cool stars

Cool dwarf stars



Science cases: Cool stars


Cool dwarf stars



High-resolution spectroscopy provides detailed information, which complements photometric observations (synergy with PLATO and ARIEL).

Atmospheric parameters of the star, in particular its chemical composition, or the presence of significant velocity fields, accretion from circumstellar material, or strong magnetic fields.




NIRSPEC spectra (Rice et al. 2010)



Strong sensitivity of potassium lines and H₂O absorption

Science cases: Cool stars

Cool evolved stars



Dynamics and **chemical composition in nearby Galaxies** (Davies et al. 2017)


Outline

- Context, the team and the process
- Sciences cases: Cool stars
- Sciences cases: Dynamical and chemical composition of stellar atmospheres
- Sciences cases: Primitive stars
- Required TRLs

Science cases: Dynamical and chemical composition of stellar atmospheres

Bergemann et al. 2019

Synthetic Sun (Staggered Sun, Miglio et al. 2013)





Stellar surface inhomogeneities (caused by convection, magnetic fields, rotation, dust...) changes with time and affect line depth, position, width.

Science cases: Dynamical and chemical composition of stellar atmospheres

Bergemann et al. 2019

Synthetic Sun (Stagger-code, Magic et al. 2013)




Stellar surface inhomogeneities (caused by convection, magnetic fields, rotation, dust...) changes with time and affect line depth, position, width.

Science cases: Dynamical and chemical composition of stellar atmospheres

Extreme-precision studies of spectral line shapes and elemental and isotopic abundances (from ~ 0.1 dex to under 0.01 dex) :

- to reveal secular changes in surface composition of stars due to diffusion, mixing, or the accretion of interstellar or protoplanetary material onto the star (Melendez et al. 2009)
- of heavy elements ($Z > 30$) in the blue (< 430 nm) to understand the neutron-capture nucleosynthesis processes, and identifying their astrophysical site(s) (Snedden et al. 2008).

High Hand in hand, the application of state-of-the-art 3D hydrodynamical model atmospheres and non-LTE modelling (e.g. Nordlander et al. 2017)




Outline

- Context, the team and the process
- Sciences cases: Cool stars
- Sciences cases: Dynamical and chemical composition of stellar atmospheres
- Sciences cases: Primitive stars
- Required TRLs

Science cases: Primitive stars

Sketch of the chemical enrichment of the Universe (Jacobson and Frebel 2014)



Understanding the nature of the very **first generation of stars**, determine the yields of early supernovae, and to **constrain the lowest metallicity** at which the gas can collapse to form low-mass stars.


Science cases: Primitive stars

Detailed elemental abundances for dozens of elements to **constrain models of supernovae** and their progenitors.

Synergy with ongoing and planned surveys (SDSS, LAMOST, MOONS, DESI, WEAVE and 4MOST) that will identify targets down to $V \sim 20$, too faint to be followed-up at higher resolution with 10m-class telescope.

The blue (and near-UV) will be crucial, as this is where the atoms have their strongest (and only visible) transitions.

Most metal poor stars known (Keller et al. 2014)



Decreasing
metallicity



Outline

- Context, the team and the process
- Sciences cases: Cool stars
- Sciences cases: Dynamical and chemical composition of stellar atmospheres

Sciences cases: Pr

- Required TRLs

Science cases: Required TRLs

Science Case	Internal rank				Wavelength range (microns)						Spectral resolution (x 1,000)						Wavelength accuracy					Polarim.	Multiplex.			Backgr. Sub.		AO		IFU		
					0.34-0.37	0.37-0.4	0.4-0.50	0.50-0.67	0.67-0.8	0.8-0.9	0.9-1.3	1.5-1.8	2.0-2.4	5	10	20	50	80	100	150	2m/s		1m/s	10cm/s	2cm/s	1cm/s	2	5	10		SCAO	LTAO
WG2: Stars and Stellar Populations																																
Cool stars	1	9	10	7																												
Primitive stars	1	7	7	7																												
Stellar atmosph.	1	7	8	6																												

= Essential

= Desirable


Wavelength range (microns)								
0.34-0.37	0.37-0.4	0.4-0.50	0.50-0.67	0.67-0.8	0.8-0.9	0.9-1.3	1.5-1.8	2.0-2.4


The red-infrared is vital for cool stars, while the blue is inescapably needed for studying the most metal-poor stars.

Wavelength stability at levels of < 0.01 km/s on time scales of a few nights, as needed to ensure the accurate studies of stellar dynamics and abundances

Science cases: Required TRLs

Science Case	Internal rank				Wavelength range (microns)							Spectral resolution (x 1,000)						Wavelength accuracy					Polarim.	Multiplex.			Backgr. Sub.	AO		IFU				
					0.34-0.37	0.37-0.4	0.4-0.50	0.50-0.67	0.67-0.8	0.8-0.9	0.9-1.3	1.5-1.8	2.0-2.4	5	10	20	50	80	100	150	2m/s	1m/s		10cm/s	2cm/s	1cm/s		2	5		10	SCAO	LTAO	
WG2: Stars and Stellar Populations																																		
Cool stars	1	9	10	7																														
Primitive stars	1	7	7	7																														
Stellar atmosph.	1	7	8	6																														

 = Essential

 = Desirable


Polarim.	Multiplex.			Backgr. Sub.
	2	5	10	


A background subtraction quality better than 0.1% in the optical and 0.5% in the infrared to avoid limiting the signal-to-noise ratio of observations

Relative flux calibration of better than 1% over spectral intervals of 20-30 nm, covering the broadest features in stellar spectra

Science cases: Required TRLs

Science Case	Internal rank				Wavelength range (microns)						Spectral resolution (x 1,000)					Wavelength accuracy					Polarim.	Multiplex.			Backgr. Sub.	AO		IFU															
					0.34-0.37	0.37-0.4	0.4-0.50	0.50-0.67	0.67-0.8	0.8-0.9	0.9-1.3	1.5-1.8	2.0-2.4	5	10	20	50	80	100	150		2m/s	1m/s	10cm/s		2cm/s	1cm/s		2	5	10	SCAO	LTAO										
WG2: Stars and Stellar Populations																																											
Cool stars	1	9	10	7																																							
Primitive stars	1	7	7	7																																							
Stellar atmosph.	1	7	8	6																																							

 = Essential

 = Desirable


AO		IFU
SCAO	LTAO	

An IFU unit enabling spatial resolution close to 10 mas. This is a requirement for resolving stellar populations in nearby galaxies and circumstellar disks.

Take home message

What can be achieved with HIRES ?

The vast light-collecting power of the ELT



It will enable **detailed** high-resolution spectroscopy of **individual stars** (faint red dwarfs and distant red giants in **nearby galaxies**)

The high-resolution of HIRES



Measuring the **absorption lines** in stellar spectra, where the thermal and turbulent velocities may be as small as 1 km/s

Measuring the **surface structures** or **anomalous chemistry**

